Higher Learning Lab · Free research resource · CC BY-NC 4.0

ADHD in Preschoolers: A Review of the Research

Bjorn, Genevive (Higher Learning Lab); Ma, Clara Fangfang (Higher Learning Lab)

Published 2025-10-28 by Higher Learning Lab, a 501(c)(3) nonprofit, Dover, Delaware.

Canonical version: https://www.higherlearninglab.org/portfolio/adhd-in-preschoolers-a-review-of-the-research/

Licensed under Creative Commons Attribution-NonCommercial 4.0.

By Higher Learning Lab

Citation: Bjorn, G., & Ma, C. F. (2025). *ADHD in preschoolers: A review of the research*. Higher Learning Lab. https://doi.org/10.5281/zenodo.23541021

Effects of Trauma and Abuse on ADHD Symptoms

Early childhood trauma and abuse greatly affect the development and expression of ADHD symptoms in young children. Research consistently indicates that children with ADHD are more likely to have encountered adverse childhood experiences (ACEs) than their peers. In a nationally representative U.S. survey of more than 76,000 children, those with ADHD had a higher prevalence of every ACE measured, and a graded relationship existed between the number of ACEs and ADHD severity: children with two, three, or four or more ACEs were significantly more likely to have moderate-to-severe ADHD (Brown et al., 2017). Specific adversities—socioeconomic hardship, parental divorce, familial mental illness, neighborhood violence, and incarceration of a household member—were each associated with higher odds of ADHD (Brown et al., 2017). These findings suggest that early psychosocial trauma can not only trigger but also exacerbate ADHD-related behaviors.

This review is meant for educational and informational purposes only. This review expressly does not give any medical advice. Always consult a qualified medical provider with medical questions.

Early childhood trauma exposure can alter the clinical presentation of ADHD. Symptoms of post-traumatic stress, such as hyperarousal, concentration difficulties, and impulsive outbursts, closely resemble core ADHD symptoms, leading to potential misdiagnosis (Szymanski et al., 2011; Weinstein et al., 2000). Children who have experienced abuse or neglect may be mistakenly diagnosed with primary ADHD when their behaviors are actually rooted in trauma-related distress (Weinstein et al., 2000). Conversely, a child with ADHD who has also experienced trauma may exhibit increased emotional dysregulation and heightened anxiety, complicating both diagnosis and treatment. Without proper screening, clinicians may overlook trauma as a significant factor contributing to a child's inattentive or hyperactive behaviors, reinforcing the need for comprehensive trauma evaluations in children suspected of having ADHD (Szymanski et al., 2011).

From a neurological perspective, chronic early stress affects brain development in ways that align with ADHD. Prolonged exposure to trauma can disrupt the hypothalamic–pituitary–adrenal (HPA) axis and raise cortisol levels, potentially hindering self-regulation and attention networks in the prefrontal cortex (Teicher & Samson, 2016). Research has shown that cumulative childhood adversity can lead to lasting changes in neural circuitry through stress-related and epigenetic pathways (McLaughlin et al., 2019). Neuroimaging studies reveal that maltreated children display atypical development in brain areas responsible for executive function and emotion regulation, which can manifest behaviorally as inattention, impulsivity, and hyperactivity (McLaughlin et al., 2019; Teicher & Samson, 2016). Given the strong link between ACEs and altered neural pathways, these findings imply that early-life stress may not only heighten the risk of ADHD but also contribute to increased symptom severity in affected children.

In summary, early trauma and abuse are not only more prevalent in children with ADHD, but they also worsen symptom severity and complicate clinical diagnosis. Consequently, clinicians highlight the critical need for trauma screening when assessing young children for ADHD (Szymanski et al., 2011). Addressing trauma through trauma-informed care—which includes therapy for PTSD symptoms, stable caregiving environments, and holistic behavioral interventions—is a vital aspect of effective ADHD management in young children exposed to adversity (Brown et al., 2017).

Sleep Disturbances and ADHD in Young Children

Difficulty Sleeping and Sleep Apnea in Young Children with ADHD

Sleep disturbances are frequently reported in preschoolers with ADHD, and research highlights a strong interplay between sleep problems and ADHD symptoms. Children with ADHD often experience bedtime resistance, delayed sleep onset, and frequent night awakenings, which can lead to fragmented sleep and difficulty waking in the morning (Beebe, 2011; Cortese et al., 2009). Objective sleep studies corroborate these parental reports, showing that young children with ADHD take longer to fall asleep, have lower sleep efficiency, and experience more sleep-disordered breathing events, such as obstructive sleep apnea (OSA) (Cortese et al., 2009). Additionally, restless movements during sleep, such as periodic limb movements or restless legs syndrome, contribute to daytime

sleepiness, further exacerbating ADHD symptoms (Cortese et al., 2009).

The Bidirectional Relationship Between Sleep and ADHD

Sleep and ADHD have a bidirectional relationship: poor sleep can worsen ADHD symptoms, and ADHD itself (or its treatments) can lead to sleep disturbances. Insufficient or fragmented sleep often results in irritability, hyperactivity, and reduced attention span the following day, behaviors that mirror or aggravate ADHD (Beebe, 2011). A longitudinal study of nearly 1,500 children found that those with persistently short sleep in the preschool years—especially before 41 months of age—displayed significantly more hyperactivity-impulsivity and lower cognitive performance at school entry (Touchette et al., 2007). Follow-up of the same cohort links childhood sleep trajectories to mental-health symptoms, including ADHD symptoms, at age 15 (Touchette et al., 2024).

Physiologically, chronic sleep deprivation and irregular sleep schedules impair attention networks and executive functioning in the developing brain, worsening core ADHD symptoms such as impulsivity and distractibility. Conversely, factors related to ADHD, including racing thoughts or high activity levels, can hinder children's ability to relax and fall asleep (Kirov & Brand, 2014). Stimulant medications, commonly prescribed for ADHD, can further complicate these challenges by increasing sleep onset latency and reducing total sleep duration (Kirov & Brand, 2014). This cycle—where ADHD worsens sleep and poor sleep exacerbates ADHD symptoms—requires clinicians to assess sleep quality in any young child with ADHD and address sleep disturbances as part of a comprehensive treatment plan (Cortese et al., 2009).

Obstructive Sleep Apnea (OSA) and ADHD

One specific sleep disorder of concern is obstructive sleep apnea (OSA), which can present with ADHD-like symptoms in children. OSA involves intermittent breathing pauses during sleep (often due to enlarged tonsils or adenoids), leading to fragmented sleep, daytime fatigue, inattention, hyperactivity, and mood disturbances. A meta-analysis of 18 studies found a medium-sized association between sleep-disordered breathing and ADHD symptoms (Hedges' $g = 0.57$), and the authors recommend screening for sleep-disordered breathing in any child with ADHD symptoms before medicating (Sedky et al., 2014).

Although pediatric OSA itself is relatively uncommon (affecting 1–4% of young children), when it does occur, it can be mistaken for or coexist with ADHD (Marcus et al., 2012). Children with ADHD are also more prone to habitual snoring and mild airway obstruction at night, even if they do not meet full OSA criteria (Chervin et al., 2002). This overlap makes careful assessment crucial; for instance, if a preschooler exhibits snoring and daytime ADHD symptoms, an evaluation for sleep apnea is warranted.

This review is meant for educational and informational purposes only. This review expressly does not give any medical advice. Always consult a qualified medical provider with medical questions.

Telltale signs of OSA include:

- Loud snoring, gasping, or pausing in breathing during sleep
- Restless sleep
- Mouth breathing, morning headaches, or excessive daytime sleepiness

Distinguishing OSA-related attention problems from primary ADHD is critical, as their treatments differ. Encouragingly, research indicates that when a child's ADHD-like symptoms are related to sleep apnea, addressing the apnea can lead to dramatic symptom improvement. Adenotonsillectomy, the surgical removal of enlarged tonsils and adenoids, is a standard treatment for pediatric OSA. A meta-analysis of 12 studies (N = 529) found a medium-sized reduction in ADHD symptoms 2–13 months after adenotonsillectomy (Hedges' g = 0.43) (Sedky et al., 2014). In a treatment-outcome study of children with both ADHD and mild OSA, adenotonsillectomy produced improvements in attention, behavior, and quality of life comparable to or greater than those seen with methylphenidate (Huang et al., 2007), and children with sleep-disordered breathing show marked decreases in hyperactivity and inattention in the year after surgery (Chervin et al., 2006). The authors of these studies recommend treating comorbid sleep-disordered breathing before medicating ADHD symptoms (Sedky et al., 2014).

Central Sleep Apnea (CSA) in Young Children

While OSA is the most well-known sleep-related breathing disorder associated with ADHD, central sleep apnea (CSA) can persist in some children even after adenotonsillectomy, particularly in those with neurodevelopmental conditions such as ADHD. Unlike OSA, CSA is caused by reduced respiratory drive from the brainstem rather than airway obstruction. Children with CSA experience pauses in breathing due to inconsistent signaling from the central nervous

system, leading to fragmented sleep, excessive daytime sleepiness, and worsened ADHD symptoms.

The brainstem, specifically the medulla oblongata and pons, plays a crucial role in regulating breathing. In children with CSA, these structures fail to send consistent signals to respiratory muscles, disrupting normal breathing patterns. The primary muscles affected include:

- **Diaphragm:** The main breathing muscle, which experiences reduced activation due to impaired brainstem signaling.
- **Intercostal muscles**: These expand and contract the chest during respiration but may not receive consistent neural input in CSA cases.
- **Upper airway muscles:** While CSA is not caused by airway obstruction, poor coordination of upper airway muscles can still disrupt breathing patterns.

Treatments for CSA in Children with ADHD

Effective treatments for CSA depend on its severity and underlying causes. Pediatric pulmonologists and sleep medicine specialists typically guide treatment. Common interventions include:

- **Supplemental oxygen therapy:** Helps maintain adequate oxygen levels and reduces apnea-related awakenings.
- **Positive airway pressure (PAP) therapy:** CPAP or BiPAP can assist children experiencing significant CSA.
- **Pharmacological interventions:** Medications like acetazolamide, a carbonic anhydrase inhibitor, may improve respiratory drive in certain CSA cases.
- **Neurological and behavioral approaches:** Addressing underlying neurological or developmental factors through behavioral therapy, structured sleep hygiene, and ADHD management may indirectly alleviate CSA symptoms.

Given the complex relationship between ADHD and sleep disorders, clinicians should assess children with ADHD for underlying sleep disturbances, especially OSA and CSA, when symptoms persist. Identifying and treating sleep-related breathing issues can significantly improve ADHD symptom management, enhance daytime functioning, and reduce the reliance on stimulant medications. Early intervention through behavioral sleep strategies, medical treatments, and evaluations for sleep disorders is essential for optimizing developmental outcomes in children with ADHD.

Improving Sleep Quality and ADHD Management

Improving sleep quality generally has positive effects on ADHD management. Behavioral sleep interventions—including establishing consistent bedtime routines, teaching children self-soothing strategies, and educating parents on sleep hygiene—have shown beneficial outcomes (Hiscock et al., 2015). In a randomized controlled trial, a brief behavioral sleep program (two sessions focusing on bedtime routines and limit-setting) for young school-age children with ADHD led to modest but significant improvements in ADHD symptoms and better sleep patterns (Hiscock et al., 2015). By three months after the intervention, parents of the treatment group reported fewer ADHD symptoms and much fewer sleep problems than the control group (30% of intervention children still had moderate-to-severe sleep problems, versus 56% in controls) (Hiscock et al., 2015). Teachers also noted improved behavior in the classroom following the sleep-focused intervention (Hiscock et al., 2015). Notably, improvements in sleep accounted for a substantial portion of the ADHD symptom reduction, indicating that better sleep translated into better self-regulation and attention during the day (Hiscock et al., 2015). These gains, while moderate, persisted at the six-month follow-up, suggesting that even short-term sleep coaching can have lasting benefits (Hiscock et al., 2015). A later translational trial delivered by community pediatricians and psychologists replicated the sleep benefits but did not find improvements in ADHD symptoms, underscoring that results depend on intervention dose and clinician supervision (Hiscock et al., 2019).

This review is meant for educational and informational purposes only. This review expressly does not give any medical advice. Always consult a qualified medical provider with medical questions.

Beyond formal interventions, simple steps like maintaining a strict bedtime, reducing screen time before bed, and ensuring the child gets adequate sleep for their age can help lessen hyperactive and impulsive behavior. In summary, sleep disturbances can both exacerbate ADHD symptoms and be a possible root cause for attention problems in some children. Identifying and treating issues like insomnia or sleep apnea in preschoolers with ADHD is an essential component of care—it can improve daytime behavior, enhance the child's mood and readiness to learn, and overall improve quality of life for both the child and their family (Cortese et al., 2009).

Supportive Interventions: Therapy, School Strategies, and Parenting Approaches

Managing ADHD in preschool-aged children typically requires a multi-faceted, supportive approach that extends beyond medication. Evidence-based therapies, educational interventions, and positive parenting strategies can effectively reduce symptoms and enhance a young child's functioning. Clinical guidelines unanimously recommend behavioral therapy as the first-line treatment for ADHD in children under six, reserving medication for cases where behavioral interventions alone are insufficient (Wolraich et al., 2019). This section outlines key supportive strategies and their effectiveness for preschoolers with ADHD.

Behavioral Therapy and Parent Instruction

The most effective and well-researched intervention for this age group is behavioral parent training. Programs that coach parents in behavior management techniques (sometimes alongside therapist-child sessions) have significantly improved young children's ADHD and oppositional behaviors (Charach et al., 2013; Rimestad et al., 2019). In these programs, therapists collaborate with parents to teach skills such as using positive reinforcement, setting consistent rules and routines, and applying appropriate, calm discipline when necessary (Centers for Disease Control and Prevention, 2021). Parents learn to anticipate and prevent problem behaviors while reinforcing desirable actions (e.g., praising the child for playing quietly for a few minutes or using token reward systems for following directions). This training also helps parents establish developmentally appropriate expectations and strengthen the parent–child relationship (Wolraich et al., 2019).

Parent training improves the child's behavior at home, reduces family stress, and provides greater stability for the child. Multiple studies confirm that parent training lessens preschoolers' inattentiveness and hyperactive behavior as reported by parents, with improvements maintained at follow-up (Rimestad et al., 2019). For example, Parent–Child Interaction Therapy (PCIT) is an evidence-based program for children under seven that utilizes live parent coaching to develop positive interaction skills and enhance compliance. It has been shown to reduce ADHD and disruptive behaviors in young children. The American Academy of Pediatrics (AAP) and CDC emphasize that only therapy focused on parental training has demonstrated efficacy in this young age group (CDC, 2021).

This review is meant for educational and informational purposes only. This review expressly does not give any medical advice. Always consult a qualified medical provider with medical questions.

Children aged 3–5 are often too immature for traditional talk therapy or cognitive-behavioral therapy (CBT) aimed at the child; instead, the parent serves as the agent of change. By learning behavior management strategies, parents can establish a more structured environment that fosters their child's success. Group-based parent training programs (such as Incredible Years or Triple P) and individualized programs (like PCIT or New Forest Parenting Programme) have both demonstrated positive outcomes. However, improvements are usually most noticeable at home and in parent reports (Rimestad et al., 2019).

Overall, behavioral therapy for preschool ADHD is associated with better attention spans, less impulsivity, and improved compliance, without the side effects medications can bring. Given this strong evidence, current guidelines strongly recommend that clinicians prescribe evidence-based parent training and behavioral therapy for preschool-aged children with ADHD before considering any medications (Wolraich et al., 2019).

School and Early Education Interventions

Even at the preschool or daycare level, structured interventions can help manage ADHD symptoms and support learning. Behavioral classroom interventions are recommended alongside parent training whenever possible (Wolraich et al., 2019). For a 3- to 5-year-old, this might involve the teacher implementing a daily behavior plan or using simple reward systems at school. Techniques such as sticker charts for good sitting or listening, very short work periods with movement breaks in between, and clear visual schedules can significantly help keep a young child with ADHD on track. Classroom management strategies shown to assist older children with ADHD (like token economies, time-outs for aggressive behavior, and frequent feedback) can be adapted for preschool settings. Small class sizes or a one-on-one aide can also offer additional supervision and prompts to help a child with ADHD thrive in an early childhood education environment. In some cases, preschoolers with severe ADHD may qualify for specialized services or an Individualized Education Program (IEP) even before kindergarten. Behavioral specialists might work with the child in preschool to practice skills like waiting their turn, following directions, and engaging in cooperative play.

Research indicates that when teachers consistently use positive reinforcement and structured routines, young children with ADHD demonstrate fewer disruptive behaviors and show greater engagement in learning tasks (Charach et al., 2013). Communication between parents and teachers is crucial—for instance, using a simple daily report or sticker chart that circulates back and forth can reinforce the same goals at home and school. Overall, early childhood educational interventions create an ADHD-friendly environment that can significantly enhance a child's self-control and social skills, setting the stage for a smoother transition to kindergarten.

Supportive Parenting Approaches

In addition to formal parent training programs, some everyday parenting techniques are especially beneficial for children with ADHD. Consistency and structure are essential. Parents should establish predictable routines—such as set times for meals, play, and bedtime—because children with ADHD thrive on consistency and clear expectations. Providing structure and clear expectations can minimize confusion and decrease opportunities for misbehavior. Positive reinforcement is a powerful tool: recognizing the child for good behavior and offering labeled praise (e.g., "I love how quietly you are coloring!") or small rewards tends to encourage those desirable behaviors (CDC, 2021). Utilizing checklists or picture charts for routines (like getting dressed or cleaning up toys) can help a young child remain focused with less frustration. Another strategy is implementing immediate and mild consequences for dangerous or destructive behavior while avoiding harsh punishment, which is often ineffective and may worsen oppositional behavior. Instead, calm and consistent time-outs or temporary loss of a privilege can be used when necessary. Parents are also encouraged to choose their battles and prioritize the most important behaviors to address while providing outlets for the child's energy (e.g., outdoor playtime) throughout the day. Parent support groups or classes can reinforce these strategies and offer moral support. By employing these approaches, parents frequently observe improvements in their child's self-regulation and decreased family stress.

Occupational and Play Therapy

Some families seek extra therapies, like occupational therapy (OT) or play therapy, to support their preschooler with ADHD. Occupational therapy can help children with co-occurring fine motor delays, sensory processing issues, or trouble with self-help skills, which are common in some kids with ADHD. An occupational therapist may focus on enhancing the child's sensory modulation—providing

activities that involve deep pressure or movement to assist a child who is overly active in calming their body. They might also teach visual and motor strategies to improve the child's attention, such as using fidget tools or scheduling "sensory breaks." While OT isn't a primary treatment for managing the core symptoms of ADHD, it can strengthen a child's ability to engage in tasks and follow routines, indirectly supporting better behavior. Play therapy represents another approach used for young children facing behavioral challenges. In play therapy, a trained therapist employs play activities and toys to help the child express emotions, practice self-control, and develop social skills. However, the CDC (2021) emphasizes that neither play therapy nor traditional talk therapy has been shown to enhance attention span or reduce impulsivity in preschoolers with ADHD.

Prognosis and Long-Term Outcomes of Preschool ADHD

One critical question for parents and clinicians is whether a very young child diagnosed with ADHD will outgrow the condition or continue experiencing difficulties later in life. Long-term studies indicate that ADHD manifesting in the preschool years often persists, although the severity and specific symptoms may evolve over time. In general, ADHD is now viewed as a chronic neurodevelopmental disorder rather than a transient phase for most individuals (Faraone et al., 2015). Follow-up research on preschoolers diagnosed with ADHD shows high rates of symptom persistence into later childhood. For example, the Preschool ADHD Treatment Study (PATS), which followed children diagnosed around age 4, found that six years later, approximately 89% of the children still met the diagnostic criteria for ADHD at around age 10 (Riddle et al., 2013). Symptom ratings did decline somewhat between preschool and early elementary school—especially among girls in the sample—but for the vast majority, ADHD symptoms remained in the moderate-to-severe clinical range through age 10 (Riddle et al., 2013). Another longitudinal study reported that after four years, 85% of children with ADHD continued to have the disorder, while only about 15% went into remission (Biederman et al., 1996). Of those who remitted, half did so in childhood and half in adolescence (Biederman et al., 1996). Thus, while some children experience a substantial reduction in symptoms as they mature, the odds indicate that a preschooler with a bona fide ADHD diagnosis will continue to struggle with attention and self-control to some degree in subsequent years.

Changes in ADHD Symptoms Over Time

That being said, the nature of ADHD symptoms often evolves with development. Clinicians observe that the overt hyperactivity observed in a 4-year-old—constant running, climbing, and the inability to sit still—tends to decrease during adolescence. As children with ADHD mature, the impulsive hyperactive behaviors typically become less apparent (Faraone et al., 2006). However, inattention and executive functioning deficits often persist or even become more pronounced as academic and organizational demands increase (Faraone et al., 2006). Long-term studies have indeed found that by adolescence, many individuals with childhood ADHD exhibit reduced hyperactivity, yet difficulties in concentration, planning, and impulsivity may continue into the teenage and adult years (Faraone et al., 2006). In a 2016 national survey, 9.4% of U.S. children aged 2–17 had ever been diagnosed with ADHD, and 89% of those still currently had the condition (Danielson et al., 2018). A meta-analysis of follow-up studies estimates that about 15% of children with ADHD still meet full diagnostic criteria at age 25, but roughly 65% continue to have clinically significant, impairing symptoms when partial remission is counted (Faraone et al., 2006). Others may achieve partial remission—they no longer meet the diagnostic criteria exactly but might still exhibit some attention deficit traits or executive function weaknesses that impact their work or daily life. Only a minority—roughly one-third—appear to fully "outgrow" ADHD by adulthood in the sense of having no impairing symptoms (Faraone et al., 2006). Thus, for many individuals, ADHD is lifelong, but it can manifest in different ways over time.

Predictors of ADHD Persistence or Remission

Researchers have sought to identify predictors that distinguish children who outgrow ADHD from those who have a more persistent course. Certain factors present in the preschool years can indicate a higher likelihood that ADHD will persist. A strong family history of ADHD (i.e., a parent or close relative with ADHD) is one such risk factor, reflecting the genetic component of the disorder (Faraone & Larsson, 2019). Children who have ADHD plus additional psychiatric or developmental disorders are also more likely to have persistent symptoms. In particular, comorbid conditions such as oppositional defiant disorder (ODD), conduct problems, anxiety, or mood disorders in a young child with ADHD predict a more chronic and severe trajectory (Biederman et al., 1996). The PATS follow-up found that the presence of ODD or conduct disorder in a preschooler with ADHD was associated with a 30% higher risk of the ADHD remaining

at age 10 (Riddle et al., 2013). High levels of psychosocial adversity (which often includes trauma, as discussed earlier) are another predictor of persistence (Biederman et al., 1996). Children from very chaotic or deprived home environments tend to have more persistent and impairing ADHD, possibly because chronic stress and lack of support hinder the development of coping skills. Conversely, children with milder symptoms and no comorbidities, who receive strong support and early intervention, are more likely to show improvement by later childhood. Some studies suggest that intellectual ability and cognitive development play a role as well—higher IQ or better executive function skills might buffer a child, whereas children with global developmental delays in addition to ADHD often have a more continuous course of difficulties.

Neurological and Brain Development Findings

Neurological and brain development findings align with these clinical observations. ADHD has been associated with differences in brain structure and maturation, particularly in the prefrontal cortex and related networks that govern attention and impulse control. Longitudinal neuroimaging research at the NIMH provided insight into the brain trajectories of children with ADHD. They found that in childhood, those with ADHD have a thinner cerebral cortex in regions important for attention and self-regulation (especially the frontal areas) compared to peers (Shaw et al., 2007). Over time, some children with ADHD showed a normalization of brain development, while others did not. Specifically, children whose ADHD symptoms remitted by adolescence tended to show catch-up growth in parts of the brain. For example, normalization of cortical thickness in the right parietal cortex was observed in children with better clinical outcomes (Shaw et al., 2006). In contrast, children who had a persistent course of ADHD showed a more “fixed” pattern of cortical thinness. Notably, those with worse outcomes had significantly thinner frontal cortex from the start, and this did not normalize with age (Shaw et al., 2006). This suggests that when the neural maturation delay in ADHD is large and remains uncorrected during development, the disorder is likely to continue, whereas if the brain can compensate or catch up in key areas, the child may essentially outgrow the clinical diagnosis.

This review is meant for educational and informational purposes only. This review expressly does not give any medical advice. Always consult a qualified medical provider with medical questions.

Importance of Early Intervention

It is important to note that early intervention can significantly influence long-term outcomes, even if it does not "cure" ADHD. The presence of ADHD symptoms in preschool is associated with worse academic and social outcomes later (for example, greater risk of grade retention, difficulties in peer relationships, and injury proneness in elementary years). Still, timely support can mitigate some risks (Charach et al., 2013).

Conclusion

Most preschoolers with true ADHD are likely to continue experiencing ADHD or related challenges later in life, although some may undergo partial or full remission of symptoms as they mature. Hyperactive symptoms generally diminish with age, but attention deficit and impulsivity can remain significant issues into adolescence and adulthood. Key indicators of a persistent course include genetics (family history), adversity and trauma, greater initial symptom severity, and co-occurring behavioral or emotional disorders. While we cannot predict with certainty the future of any one child, ongoing research into brain development is illuminating why some children outgrow ADHD. The prevailing understanding is that ADHD is often a lifelong condition, but early supportive interventions and a nurturing environment can enhance the child's long-term outlook. Children who receive assistance in preschool—through behavioral therapy, educational support, and family training—are better prepared to manage their symptoms, which can lead to improved academic performance and social relationships, even if ADHD does not completely disappear. Therefore, identifying and treating ADHD early in life provides the best opportunity for a child to develop effective coping strategies and to flourish as they grow, potentially changing the trajectory of the disorder for the better (Biederman et al., 1996).

Key Takeaways from the Research on ADHD in Preschoolers

1. **Trauma can increase ADHD risk and severity:** Children with ADHD are more likely to have experienced adverse childhood experiences (ACEs), such as abuse, neglect, or

exposure to violence. These experiences can both trigger and worsen ADHD symptoms.

2. **Trauma symptoms can mimic ADHD:** Symptoms of post-traumatic stress, such as hyperarousal and impulsivity, resemble ADHD symptoms, sometimes leading to misdiagnosis if trauma is not correctly assessed and healed.
3. **Brain development is impacted by stress and ADHD:** Chronic early stress can alter brain structures responsible for self-regulation and attention, making ADHD symptoms more severe. Neurological research shows that children with ADHD often have delayed cortical development in brain regions linked to impulse control and focus.
4. **Sleep problems are common in children with ADHD:** Many preschoolers with ADHD struggle with falling and staying asleep, which can worsen hyperactivity, impulsivity, and inattention. Addressing sleep problems can improve daytime behavior.
5. **Obstructive sleep apnea (OSA) can mimic ADHD:** Sleep-disordered breathing is moderately associated with ADHD symptoms, and treating it (e.g., adenotonsillectomy) produces measurable reductions in ADHD symptoms (Sedky et al., 2014).
6. **Central sleep apnea (CSA) is linked to brainstem issues:** Some children with ADHD also experience CSA, where the brain does not properly signal the body to breathe during sleep. It is uncommon and requires evaluation by a pediatric sleep specialist.
7. **Parent training is the most effective first-line treatment for ADHD in preschoolers.** Behavioral parent training programs, such as Parent-Child Interaction Therapy (PCIT), Incredible Years, and Triple P, help parents manage ADHD symptoms without medication and improve child behavior.
8. **Early intervention and structure are crucial:** Providing consistent routines, clear expectations, and positive reinforcement at home and school can help preschoolers with ADHD develop self-control and coping skills.
9. **Most preschoolers with ADHD will continue to have symptoms into later childhood:** Long-term studies show that about 85-89% of preschoolers diagnosed with ADHD still meet the criteria for the disorder at age 10, and 50-65% will continue to experience ADHD-related challenges into adulthood.
10. **ADHD symptoms can improve with the right support:** While ADHD is often a lifelong condition, early

behavioral interventions, trauma-informed care, and proper sleep management can help children develop skills to manage symptoms and succeed in school and relationships.

Action Steps for Parents and Caregivers

1. **Ensure trauma and sleep issues are evaluated**: If your child has ADHD symptoms, ask your pediatrician to **screen for trauma history and sleep disorders**, including **obstructive and central sleep apnea**. Identifying and treating these issues can improve symptoms.
2. **Participate in parent training programs**: Programs like **PCIT, Incredible Years, and Triple P** teach caregivers to **set clear expectations, use positive reinforcement, and manage challenging behaviors** without relying on medication.
3. **Create a consistent daily routine**: Structure helps children with ADHD thrive. Set **regular bed, meal, and play times** to stabilize and reduce impulsive behaviors.
4. **Improve sleep hygiene**: Limit screen time to 30 minutes per day (or less), stop screens at least two hours before bed, establish **calming bedtime routines**, and seek medical advice if your child snores, has trouble sleeping, or wakes up frequently.
5. **Work with teachers and specialists**: Advocate for **behavioral strategies in preschool**, such as **visual schedules, movement breaks, and clear expectations**. Request an **Individualized Education Program (IEP)** or specialized support services from the school district as needed.

By **understanding ADHD, addressing sleep and trauma-related factors, and using behavioral strategies**, parents and caregivers can **help their children manage symptoms, develop self-regulation skills, and succeed at home and school**.

About the Author

Genevive Bjorn, Ed.D., is an education researcher and consultant specializing in workforce development and professional development. She holds a doctorate in education from Johns Hopkins University. Before that, she was a national-award-winning secondary science teacher in Chula Vista, CA, with experience in collaborative special and general education.

References

Beebe, D. W. (2011). Cognitive, behavioral, and functional consequences of inadequate sleep in children and adolescents. Pediatric Clinics of North America, 58(3), 649–665. https://doi.org/10.1016/j.pcl.2011.03.002

Biederman, J., Faraone, S., Milberger, S., Curtis, S., Chen, L., Marrs, A., Ouellette, C., Moore, P., & Spencer, T. (1996). Predictors of persistence and remission of ADHD into adolescence: Results from a four-year prospective follow-up study. Journal of the American Academy of Child & Adolescent Psychiatry, 35(3), 343–351. https://doi.org/10.1097/00004583-199603000-00016

Brown, N. M., Brown, S. N., Briggs, R. D., Germán, M., Belamarich, P. F., & Oyeku, S. O. (2017). Associations between adverse childhood experiences and ADHD diagnosis and severity. Academic Pediatrics, 17(4), 349–355. https://doi.org/10.1016/j.acap.2016.08.013

Centers for Disease Control and Prevention. (2024). Parent training in behavior management for ADHD. https://www.cdc.gov/adhd/treatment/parent-training.html

Charach, A., Carson, P., Fox, S., Ali, M. U., Beckett, J., & Lim, C. G. (2013). Interventions for preschool children at high risk for ADHD: A comparative effectiveness review. Pediatrics, 131(5), e1584–e1604. https://doi.org/10.1542/peds.2012-0974

Chervin, R. D., Archbold, K. H., Dillon, J. E., Panahi, P., Pituch, K. J., Dahl, R. E., & Guilleminault, C. (2002). Inattention, hyperactivity, and symptoms of sleep-disordered breathing. Pediatrics, 109(3), 449–456. https://doi.org/10.1542/peds.109.3.449

Chervin, R. D., Ruzicka, D. L., Giordani, B. J., Weatherly, R. A., Dillon, J. E., Hodges, E. K., Marcus, C. L., & Guire, K. E. (2006). Sleep-disordered breathing, behavior, and cognition in children before and after adenotonsillectomy. Pediatrics, 117(4), e769–e778. https://doi.org/10.1542/peds.2005-1837

Cortese, S., Faraone, S. V., Konofal, E., & Lecendreux, M. (2009). Sleep in children with attention-deficit/hyperactivity disorder: Meta-analysis of subjective and objective studies. Journal of the American Academy of Child & Adolescent Psychiatry, 48(9), 894–908. https://doi.org/10.1097/CHI.0b013e3181ac09c9

Danielson, M. L., Bitsko, R. H., Ghandour, R. M., Holbrook, J. R., Kogan, M. D., & Blumberg, S. J. (2018). Prevalence of parent-reported ADHD diagnosis and associated treatment among U.S. children and adolescents, 2016. Journal of Clinical Child & Adolescent Psychology, 47(2), 199–212. https://doi.org/10.1080/15374416.2017.1417860

Faraone, S. V., Asherson, P., Banaschewski, T., Biederman, J., Buitelaar, J. K., Ramos-Quiroga, J. A., Rohde, L. A., Sonuga-Barke, E. J., Tannock, R., & Franke, B. (2015). Attention-deficit/hyperactivity disorder. Nature Reviews Disease Primers, 1, 15020. https://doi.org/10.1038/nrdp.2015.20

Faraone, S. V., Biederman, J., & Mick, E. (2006). The age-dependent decline of attention deficit hyperactivity disorder: A meta-analysis of follow-up studies. Psychological Medicine, 36(2), 159–165. https://doi.org/10.1017/S003329170500471X

Faraone, S. V., & Larsson, H. (2019). Genetics of attention deficit hyperactivity disorder. Molecular Psychiatry, 24(4), 562–575. https://doi.org/10.1038/s41380-018-0070-0

Hiscock, H., Mulraney, M., Heussler, H., Rinehart, N., Schuster, T., Grobler, A. C., ... & Sciberras, E. (2019). Impact of a behavioral intervention, delivered by pediatricians or psychologists, on sleep problems in children with ADHD: A cluster‐randomized, translational trial. *Journal of Child Psychology and Psychiatry*, *60*(11), 1230-1241. https://doi.org/10.1111/jcpp.13083

Hiscock, H., Sciberras, E., Mensah, F., Gerner, B., Efron, D., Khano, S., & Oberklaid, F. (2015). Impact of a behavioural sleep intervention on symptoms and sleep in children with attention deficit hyperactivity disorder, and parental mental health: Randomised controlled trial. BMJ, 350, h68. https://doi.org/10.1136/bmj.h68

Huang, Y.-S., Guilleminault, C., Li, H.-Y., Yang, C.-M., Wu, Y.-Y., & Chen, N.-H. (2007). Attention-deficit/hyperactivity disorder with obstructive sleep apnea: A treatment outcome study. Sleep Medicine, 8(1), 18–30. https://doi.org/10.1016/j.sleep.2006.05.016

Kirov, R., & Brand, S. (2014). Sleep problems and their effect in ADHD. Expert Review of Neurotherapeutics, 14(3), 287–299. https://doi.org/10.1586/14737175.2014.885382

Marcus, C. L., Brooks, L. J., Draper, K. A., Gozal, D., Halbower, A. C., Jones, J., Schechter, M. S., Sheldon, S. H., Spruyt, K., Ward, S. D., Lehmann, C., & Shiffman, R. N. (2012). Diagnosis and management of childhood obstructive sleep apnea syndrome. Pediatrics, 130(3), 576–584. https://doi.org/10.1542/peds.2012-1671

McLaughlin, K. A., Weissman, D., & Bitrán, D. (2019). Childhood adversity and neural development: A systematic review. Annual Review of Developmental Psychology, 1, 277–312. https://doi.org/10.1146/annurev-devpsych-121318-084950

Riddle, M. A., Yershova, K., Lazzaretto, D., Paykina, N., Yenokyan, G., Greenhill, L., Abikoff, H., Vitiello, B., Wigal, T., McCracken, J. T., Kollins, S. H., Murray, D. W., Wigal, S., Kastelic, E., McGough, J. J., dosReis, S., Bauzó-Rosario, A., Stehli, A., & Posner, K. (2013). The Preschool Attention-Deficit/Hyperactivity Disorder Treatment Study (PATS) 6-year follow-up. Journal of the American Academy of Child & Adolescent Psychiatry, 52(3), 264–278.e2. https://doi.org/10.1016/j.jaac.2012.12.007

Rimestad, M. L., Lambek, R., Zacher Christiansen, H., & Hougaard, E. (2019). Short- and long-term effects of parent training for preschool children with or at risk of ADHD: A systematic review and meta-analysis. Journal of Attention Disorders, 23(5), 423–434. https://doi.org/10.1177/1087054716648775

Sankaran, D., Lakshminrusimha, S., & Manja, V. (2022). Methamphetamine: burden, mechanism and impact on pregnancy, the fetus, and newborn. *Journal of Perinatology*, *42*(3), 293-299. https://doi.org/10.1038/s41372-021-01271-8

Sedky, K., Bennett, D. S., & Carvalho, K. S. (2014). Attention deficit hyperactivity disorder and sleep disordered breathing in pediatric populations: A meta-analysis. Sleep Medicine Reviews, 18(4), 349–356. https://doi.org/10.1016/j.smrv.2013.12.003

Shaw, P., Eckstrand, K., Sharp, W., Blumenthal, J., Lerch, J. P., Greenstein, D., Clasen, L., Evans, A., Giedd, J., & Rapoport, J. L. (2007). Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation. Proceedings of the National Academy of Sciences, 104(49), 19649–19654. https://doi.org/10.1073/pnas.0707741104

Shaw, P., Lerch, J., Greenstein, D., Sharp, W., Clasen, L., Evans, A., Giedd, J., Castellanos, F. X., & Rapoport, J. (2006). Longitudinal mapping of cortical thickness and clinical outcome in children and adolescents with attention-deficit/hyperactivity disorder. Archives of General Psychiatry, 63(5), 540–549. https://doi.org/10.1001/archpsyc.63.5.540

Szymanski, K., Sapanski, L., & Conway, F. (2011). Trauma and ADHD – Association or diagnostic confusion? A clinical perspective. Journal of Infant, Child, and Adolescent Psychotherapy, 10(1), 51–59. https://doi.org/10.1080/15289168.2011.575704

Teicher, M. H., & Samson, J. A. (2016). Annual Research Review: Enduring neurobiological effects of childhood abuse and neglect. Journal of Child Psychology and Psychiatry, 57(3), 241–266. https://doi.org/10.1111/jcpp.12507

Touchette, E., Fréchette-Boilard, G., Petit, D., Geoffroy, M. C., Pennestri, M. H., Côté, S., ... & Montplaisir, J. (2024). Longitudinal study of childhood sleep trajectories and adolescent mental health problems. *Sleep Advances*, *5*(1), zpae013. https://doi.org/10.1093/sleepadvances/zpae013

Touchette, É., Petit, D., Séguin, J. R., Boivin, M., Tremblay, R. E., & Montplaisir, J. Y. (2007). Associations between sleep duration patterns and behavioral/cognitive functioning at school entry. Sleep, 30(9), 1213–1219. https://doi.org/10.1093/sleep/30.9.1213

Weinstein, D., Staffelbach, D., & Biaggio, M. (2000). Attention-deficit hyperactivity disorder and posttraumatic stress disorder: Differential diagnosis in childhood sexual abuse. Clinical Psychology Review, 20(3), 359–378. https://doi.org/10.1016/S0272-7358(98)00107-X

Wolraich, M. L., Hagan, J. F., Allan, C., Chan, E., Davison, D., Earls, M., Evans, S. W., Flinn, S. K., Froehlich, T., Frost, J., Holbrook, J. R., Lehmann, C. U., Lessin, H. R., Okechukwu, K., Pierce, K. L., Winner, J. D., & Zurhellen, W. (2019). Clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. Pediatrics, 144(4), e20192528. https://doi.org/10.1542/peds.2019-2528